

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A23L 1/304, 1/30, A61K 33/24</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/42211</b> <b>(43) International Publication Date:</b> 1 October 1998 (01.10.98)
<b>(21) International Application Number:</b> PCT/US98/05646 <b>(22) International Filing Date:</b> 24 March 1998 (24.03.98) <b>(30) Priority Data:</b> 08/822,483 24 March 1997 (24.03.97) US <b>(71) Applicant:</b> PROGRESSIVE RESEARCH LABS [US/US]; Suite 162, 9219 Katy Freeway, Houston, TX 77024 (US). <b>(72) Inventor:</b> WOMACK, Rick, W.; Suite 50, 9707 Richmond Avenue, Houston, TX 77042 (US). <b>(74) Agent:</b> STREETS, Jeffrey, L.; Patterson & Streets, L.L.P., Suite 1500, 3040 Post Oak Boulevard, Houston, TX 77056 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> NUTRITIONAL SUPPLEMENTS FOR IMPROVING GLUCOSE METABOLISM  <b>(57) Abstract</b>  The invention provides nutritional supplements and methods for administering nutritional supplements that improve glucose metabolism, particularly for persons with diabetes. A first nutritional supplement, or "Phase I" supplement, comprises a source of vanadate and a source of chromium. A second nutritional supplement, or "Phase II" supplement, comprises <i>Gymnema sylvestre</i> and lipoic acid. The nutritional supplements are alternated to prevent accumulation of the nutrients in the body and also to overcome desensitization that can occur over long periods of continuous use. While the nutritional supplements may be alternated at almost any frequency and taken over almost any duration, it is preferred that each Phase be taken for between about 2 and about 6 months, most preferably about 3 months or about 90 days, before alternating back to the other Phase.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## NUTRITIONAL SUPPLEMENTS FOR IMPROVING GLUCOSE METABOLISM

This is a continuation-in-part of application Serial No. 08/425,582 filed on April 20,  
5 1995.

### FIELD OF THE INVENTION

The present invention relates to nutritional supplements and methods of using them. More particularly, the present invention relates to a nutritional supplements which assist in the  
10 metabolism of glucose.

### BACKGROUND OF THE DISCLOSURE

Diabetes mellitus is caused in almost all instances by diminished rates of secretion of insulin by the beta cells of the islets of Langerhans in the pancreas. Diabetes is usually divided  
15 into two different types: juvenile diabetes that usually, but not always, begins in early life, and maturity-onset diabetes that usually, but not always, begins in later life and mainly in obese persons.

Maturity-onset type of diabetes is likely to occur in those with a family history of diabetes and is characterized by blurred vision, itching, unusual thirst, drowsiness, obesity,  
20 fatigue, skin infections, slow healing, and tingling or numbness in the feet. Onset of symptoms is usually later in life. The maturity-onset type of diabetes seems to result from degeneration or suppression of the beta cells as a result of more rapid aging in susceptible persons than in others.

Obesity predisposes an individual to this type of diabetes, probably for two different reasons. First the beta cells of the islets of Langerhans in an obese person become less responsive to  
25 stimulation by increased blood glucose levels. Therefore, the surge of insulin secretion following a meal is less marked in obese persons. Second, obesity also greatly decreases the number of insulin receptors in the insulin target cells throughout the body. For these reasons,

increased quantities of insulin are required to have the same metabolic effects in obese persons as in persons who are not obese.

Most of the pathology of diabetes mellitus can be attributed to one of three major effects of insulin lack. First, low levels of insulin cause a decrease in the utilization of glucose by the body cells with a resultant increase in blood glucose concentration to as high as 300 to 122 mg/dl. Second, insulin lack causes a markedly increased mobilization of fats from the fat storage areas, resulting in abnormal fat metabolism as well as deposition of lipids in vascular walls to cause atherosclerosis. Third, insulin lack can result in a depletion of protein in the tissues of the body.

Typical treatment of diabetes mellitus, including full-blown cases of maturity-onset diabetes, involves administering enough insulin so that the patient will have as nearly normal carbohydrate, fat, and protein metabolism as possible. Optimal therapy can prevent most acute effects of diabetes and greatly delay the chronic effects as well.

Insulin is available in several different forms. Regular insulin has a duration of action lasting from 3 to 8 hours, whereas other forms of insulin are absorbed slowly from the injection site and therefore have effects that last as long as 10 to 48 hours. Ordinarily, the severely diabetic patient is given a single dose of a longer-acting insulin each day to increase overall carbohydrate metabolism throughout the day. Then additional quantities of regular insulin are given at those times of the day when the blood glucose level tends to rise too high, such as at meal times. Thus, each patient is established on an individualized pattern of treatment.

Frequently, following a special diet can control maturity-onset diabetes sufficiently so that insulin is no longer required. It is recommended that an individual with maturity-onset diabetes follow a high-carbohydrate, high-fiber diet to reduce the need for insulin and lower the fat levels in the blood.

An estimated 5.5 million Americans are being treated for diabetes. In addition, studies estimate that there are 5 million adults with undetected maturity-onset diabetes and another 20

million having impaired glucose tolerance that may lead to full-blown diabetes. The National Institutes of Health report that undiagnosed diabetes is the reason behind millions losing their vision. Diabetes is the third leading cause of death in the United States.

Therefore, there is a need for a composition and method for enhancing glucose metabolism in individuals with maturity-onset diabetes and thereby reduce or prevent the necessity of using insulin. It would be desirable if this composition and method were convenient to administer and cost less than insulin so that it could be easily afforded by individuals with low incomes who have no insurance coverage. It would also be desirable if such a composition could be purchased over the counter, thereby making it more widely available to individuals at high risk of maturity-onset diabetes.

## SUMMARY OF THE INVENTION

The present invention provides a nutritional system for improving glucose metabolism, comprising a first supplement comprising a source of vanadate and a source of chromium and a second supplement comprising *Gymnema sylvestre* and lipoic acid. *Gymnema sylvestre* is provided as an extract from *Gymnema sylvestre* leaves. The preferred source of vanadate is vanadyl sulfate and the preferred source of chromium is selected from the group consisting of chromium picolinate, chromium glucose tolerance factor, and mixtures thereof. The source of chromium may comprise both chromium picolinate and chromium glucose tolerance factor. It may be desirable for the first supplement to further comprises L-carnitine.

Another aspect of the invention provides a nutritional system for improving glucose metabolism, comprising a first supplement comprising a source of vanadate and a second supplement comprising a component selected from the group consisting of *Gymnema sylvestre*, lipoic acid and combinations thereof, wherein the second supplement is substantially free from vanadate. The first supplement preferably further comprises a source of chromium.

Yet another aspect of the invention provides a method of administering a nutritional supplements. The method comprises the steps of administering a daily dosage of a first nutritional supplement over a first time period and administering a daily dosage of a second nutritional supplement over a second time period following the first time period. These steps  
5 may be repeated as desired. In one embodiment, the first nutritional supplement comprises a source of vanadate and a source of chromium, and the second nutritional supplement comprises *Gymnema sylvestre* and lipoic acid. In an alternative embodiment, the first nutritional supplement comprises *Gymnema sylvestre* and lipoic acid, and the second nutritional supplement comprises a source of vanadate and a source of chromium. The first and second time periods  
10 may be the same or different, preferably between about 2 and about 6 months, most preferably about 3 months or about 90 days.

#### **DETAILED DESCRIPTION OF A PREFERRED EMBODIMENT**

The present invention provides a nutritional supplement that enhances glucose  
15 metabolism. While the supplement may be used by individuals with no apparent symptoms of diabetes, the supplement is ideal for use by individuals with maturity-onset diabetes or juvenile diabetes to prevent, reduce or eliminate the necessity of using insulin. The supplement contains ingredients which work together to enhance the effect of insulin on the regulation of glucose concentration in the blood by facilitating or assisting metabolism of glucose in the cells of the  
20 body.

In one aspect of the invention, a nutritional supplement is provided which comprises a source of vanadate, such as vanadyl sulfate, and a source of chromium, such as chromium picolinate. The nutritional supplement produces insulin-like effects that prevent, reduce or eliminate the need to administer insulin. It is preferred that the supplement be taken throughout  
25 the day in order to make the vanadate and chromium available to the cells when needed. It is particularly preferred that the daily dosage be taken in approximately equal amounts with

breakfast, lunch and dinner in order for the nutrients to assist in glucose metabolism following mealtime when they are needed the most. Optionally, the nutritional supplement may further comprise L-carnitine.

In another aspect of the invention, an alternative nutritional supplement is provided with  
5 comprises a source of gymnema sylvestre and a source of lipoic acid. This alternative nutritional supplement also produces insulin-like effects the prevent, reduce or eliminate the need to administer insulin. It is preferred that the supplement be taken throughout the day in order to make the vanadate and chromium available to the cells when needed. It is particularly preferred that the daily dosage be taken in approximately equal amounts with breakfast, lunch and dinner  
10 in order for the nutrients to assist in glucose metabolism following mealtime when they are needed the most.

In yet another aspect of the invention, a plurality of nutritional supplements and a nutritional program are provided. The plurality of nutritional supplements comprise a "Phase I" supplement comprising a source of vanadate and a source of chromium and a "Phase II"  
15 supplement comprising a source of gymnema sylvestre and a source of lipoic acid. The plurality of nutritional supplements are alternated to prevent accumulation of the nutrients in the body and also to overcome desensitization that can occur after a nutrient is taken over a period of time. While the nutritional supplements may be alternated at almost any frequency and taken over almost any duration, it is preferred that each Phase be taken for between about 2 and about 6  
20 months, most preferably between about 2 and about 3 months, before alternating back to the other Phase.

In a further aspect of the invention, the nutritional supplement may optionally include other certain vitamins, minerals, amino acids, enzymes and/or herbs. These additional ingredients may be included in a common pill along with the vanadyl sulfate, chromium  
25 picolinate and L-carnitine or may be taken as a separate pill, preferably taken more or less simultaneously.

Nutritional supplements according to the present invention are not intended to eliminate the need for an individual to follow a program of appropriate diet and exercise, nor does it eliminate the need for insulin in all cases. Rather, the supplements of the present invention improve glucose metabolism by enhancing the effects of natural or administered levels of insulin and an appropriate diet and exercise program.

A nutritional supplement containing effective amounts of metabolically available forms of vanadate and chromium will improve glucose metabolism, particularly in individuals with maturity-onset diabetes. Vanadate and chromium perform different insulin-like functions which, when administered in appropriate ratios and forms, enhance glucose metabolism in substantially the same way as insulin itself.

As discussed above, insulin performs at least two important functions. First, insulin improves the utilization of glucose by the body cells which in turn controls blood glucose concentrations from getting too high. Second, insulin transports fatty acids to fat cells for storage, thereby reducing the mobilization of fats from fat storage areas which can result in abnormal fat metabolism as well as deposition of lipids in vascular walls to cause atherosclerosis.

The first function of insulin mentioned above, that of improving the utilization of glucose, is enhanced by the combination of vanadate and chromium. Vanadate ions, like insulin, stimulate glucose transport, activate glycogen synthase, increase glycogen syntheses in fat cells, and stimulate carbohydrate uptake in the liver. Glycogen synthase is an enzyme which causes the conversion of glucose into glycogen. Glycogen itself is a polysaccharide which is the chief carbohydrate storage material in humans. It has been found that maximum glycogen synthase activation produced by vanadate is indistinguishable from that of insulin. Evidence that strongly suggests a common mechanism of action for insulin and vanadate includes the following findings: (1) with maximum insulin, additional quantities of vanadate are without effect; (2) with sub maximal insulin, additional quantities of vanadate increase both the glycogen synthase activation state and 2-deoxyglucose transport to the level obtained with maximum insulin; (3)



insulin and vanadate counteract the activating effect of adrenaline on glycogen phosphorylase in a similar manner; adrenaline partially reverses vanadate and insulin activate glycogen synthase within similar time frames. Thus, the presence of invivo vanadate can lead to improved glucose metabolism and enhance the effects of natural or administered levels of insulin.

5           The preferred source of vanadate is vanadyl sulfate. Once ingested, vanadyl sulfate typically forms vanadate which is a salt of vanadic acid. While doses as low as 10 mg of vanadyl sulfate may provide some enhancement of glucose metabolism, the preferred dosage of vanadyl sulfate for an individual with diabetes weighing from about 150 pounds to about 250 pounds is in the range between about 30 mg and 150 mg per day. The most preferred dosage of vanadyl  
10 sulfate is about 60 mg per day. Whereas vanadyl sulfate has been previously included in certain dietary supplements, dosages ranging between 1 mcg and 1,000 mcg (1 mg) are inadequate to provide significant insulin-like effects.

          Chromium, like vanadyl sulfate, possesses properties which both mimic and enhance the effects of insulin. Chromium enhances the effects of insulin by indirectly assisting amino acid  
15 uptake by muscles, stimulates protein synthesis and retards the rate of protein breakdown. Many clinical studies with supplemental chromium have shown only modest improvements in glucose tolerance due to poor absorption of nutritional (trivalent) chromium. In this respect, trivalent chromium has a strongly positive charge that impedes its movement across cell membranes. Due to the presence of competing ions such as copper, iron, manganese and zinc in the human body,  
20 adequate absorption of chromium occurs only when the metal is associated with a natural chelating agent, such as picolinic acid. Because of its unique structure, picolinic acid has a strong affinity for transitional metals such as zinc, manganese, and chromium. In this respect, it binds tightly to these metals thereby neutralizing their positive charges and expediting their movement across cell membranes.

25           It is preferred that chromium be provided in a biologically active form of chromium, particularly chromium picolinate or chromium glucose tolerance factor. In chromium picolinate,

the picolinic acid serves as an effective metal chelator that improves the utilization and uptake of the chromium and plays an important physiological roll in trace mineral absorption. Chromium is believed to be the active factor while the picolinic acid serves as the chelator to improve bioavailability. Animal studies have shown chromium picolinate to be absorbed and retained  
5 five to ten times better than other forms of chromium and have also found it to be remarkable safe. The preferred dosage of chromium picolinate for an individual with diabetes weighing from about 150 pounds to about 250 pounds is in the range between about 150 and about 600 mcg per day, with about 300 mcg per day being most preferred. The preferred dosage of chromium glucose tolerance factor is between about 50 and about 400 mcg per day, with about  
10 100 mcg per day being most preferred. While the supplement may include only one form of chromium, it is preferred that the supplement include both forms in their preferred doses given above.

The second function of insulin mentioned above, that of transporting fatty acids to fat cells for storage, may be enhanced by L-carnitine. By preventing fatty build-up, this amino acid  
15 aids in weight loss, decreases the risk of heart disease, and improves athletic ability. Carnitine can be manufactured in the body if sufficient amounts of lysine, B1, B6 and iron are available. However, vegetarians are more likely to be deficient in carnitine due to a diet that is low in lysine. The preferred dosage of L-carnitine for an individual with diabetes weighing from about 150 to about 250 pounds is in the range between about 100 mg and about 1,000 mg per day. The  
20 most preferred dosage of L-carnitine is about 150 mg.

Because L-carnitine can be manufactured in the body, it is possible to complement the L-carnitine with sufficient amounts of lysine, vitamin B1, vitamin B6 and iron to facilitate production of L-carnitine. Therefore, while lysine, vitamin B1, vitamin B6 and iron are not essential to the function of the present invention, it is preferred that the supplement include  
25 between about 5 and about 10 mg per day of both vitamins B1 and B6, and between about 10 mg and about 25 mg of iron. It is most preferred that vitamins B1 and B6 be supplied by a dose of

between about 30 mg and about 70 mg of a B-complex.

Two water soluble extracts, GS3 and GS4, obtained from the leaves of *Gymnema sylvestre*, a woody climber growing in the tropical forests of central and southern India, may be used to bring about blood glucose homeostasis through increased serum insulin levels. It is  
5 believed that *Gymnema sylvestre* appears to enhance endogenous insulin, possibly by regeneration/revitalization of residual beta cells in the endocrine pancreas that are responsible for insulin production. Daily supplementation with *Gymnema sylvestre* has been shown to cause a significant reduction in blood glucose, glycosylated hemoglobin and glycosylated plasma proteins, thereby allowing conventional drug dosages to be decreased. Both juvenile and adult  
10 onset diabetes appear to respond to the action of *Gymnema sylvestre*.

Lipoic acid has been found to exhibit antioxidant properties and metabolic enhancement which facilitates regeneration of damaged nerves. Lipoic acid also appears to reduce the degree of glycation, or reaction, of proteins caused by excess blood sugar, which is common in  
15 diabetics. It is generally recognized that many of the metabolic complications that occur in diabetics are a result of persistent elevation of blood sugar, which then attaches to the blood proteins. Lipoic acid substantially reduces glycation, increases insulin sensitivity, and may lower blood sugar levels.

The present invention provides a first nutritional supplement for diabetics which combines vanadyl sulfate and chromium picolinate. It is also preferred that the first nutritional  
20 supplement include L-carnitine. The present invention also provides a second nutritional supplement for diabetics which combines *Gymnema sylvestre* and lipoic acid.

The supplements of the present invention are formulated to produce insulin-like effects that prevent, reduce or eliminate the need to administer insulin. It is preferred that either supplement be taken throughout the day to be available to the cells as needed. It is particularly  
25 preferred that the daily dosage be taken in approximately equal amounts with breakfast, lunch and dinner. Each of the supplements of the present invention provide nutrients to assist in

glucose metabolism following mealtime when they are needed the most.

The present invention further provides a nutritional system and method for diabetics which avoids desensitization that can occur after taking the same supplement continuously for a prolonged period of time. The nutritional system comprises both the first and second nutritional supplements, described above, maintained in separate capsules, preferably even in separate bottles. The nutritional system is administered by taking a daily dosage of either the first or second nutritional supplements for a period of months, then switching to the other nutritional supplement for a period of months. The period of months for administering the first and second nutritional supplements may be the same or different lengths of time. However, the preferred period of months is between about 2 and about 6 months, with the even more preferred period of months being between about 2 and about 4 months. The most preferred period of months is about 3 months or 90 days.

It is noted that the first and second nutritional supplements could be combined in a single pill, alternated between meals, alternated daily or alternated at any other frequency, and still be effective for reducing the effects of insulin lack. However, the cells of the body would eventually become desensitized to such supplements or methods of administering the supplements because of the substantially continuous presence of the nutrients. The present invention is most effective over extended period of time by alternating supplements after a period of months. It is important to understand that the period of months allows sufficient time for the body to clear itself of trace amounts of the nutrients and to avoid desensitization. Accordingly, as other nutrients are found to have beneficial effects on glucose metabolism, it may be desirable to provide additional nutritional supplements that can be administered in a rotation. In this manner, the period of months between successive use of the same supplement could be extended.

It may be desirable that the nutritional supplements include other certain vitamins, minerals, amino acids, enzymes and/or herbs. These additional ingredients may be included in a

single pill along with the vanadyl sulfate, chromium picolinate and L-carnitine or may be taken as a separate pill taken more or less simultaneously.

Vitamins that are particularly beneficial to the metabolism of glucose are the B-complex vitamins and vitamin A. Even where the supplement includes L-carnitine, it is preferred that the supplement include between about 5 and about 10 mg per day of both vitamins B1 and B6, and between about 40 mg and about 60 mg per day of niacinamide (vitamin B3). It is most preferred that the supplement include a dose of between about 30 mg and about 70 mg per day of B-complex vitamins where the amounts of vitamins B1, B3 and B6 included therein are about 6 mg, 50 mg and 6 mg, respectively.

Vitamin A is necessary to the utilization of protein. It may be taken as beta-carotene, which is converted to vitamin A in the liver, or as the vitamin itself. The preferred dose of vitamin A is between about 10,000 IU and about 20,000 IU per day, with the most preferred dose being about 15,000 IU per day.

Minerals are also important in the metabolism of glucose and may be ingested from food or included in a supplement. In order to be certain that sufficient amounts of the most important minerals are available, it is preferred that the supplement also include magnesium, potassium, calcium, copper, selenium, and zinc. The preferred daily doses of these minerals is about 380 mg magnesium, about 114 mg potassium, about 760 mg calcium, about 2 mg copper, about 80 mcg selenium, and about 50 mg zinc. It should be recognized that the amounts of these minerals can vary widely within the scope of the present invention.

L-glutamine is an important amino acid in curbing fatigue and the craving for sugar. The preferred dose of L-glutamine is between about 100 mg and about 500 mg per day, with about 300 mg per day being most preferred.

Many different enzymes may be incorporated into the supplement, in accordance with the invention, to assist in the digestion of food. The preferred enzymes include pancreatin, amylase, papain, lipase and betaine.

Pancreatin is an enzyme derived from the secretions of an animal pancreas. It is preferred that the supplement include pancreatin in doses ranging between about 50 and about 150 mg per day. It is most preferred that the supplement include about 100 mg pancreatin per day.

5 Amylase is an effective digestive enzyme secreted in high concentrations in the human body. Amylase is found in saliva and works to break down carbohydrates. It is preferred that the supplement include amylase in doses ranging between about 50 and about 150 mg per day. It is most preferred that the supplement include about 100 mg amylase per day.

Papain is a proteolytic enzyme that works exclusively to break down proteins. It is preferred that the supplement include papain in doses ranging between about 25 mg and about 70  
10 mg per day each. It is most preferred that the supplement include about 45 mg per day of papain.

Lipase is an enzyme that aids in the digestion of fat. It is preferred that the supplement include lipase in doses ranging between about 80 mg and about 230 mg per day each. It is most preferred that the supplement include about 150 mg lipase per day.

Betaine, a preferred form of hydrochloric acid (HCl), aids in the digestion of tough  
15 foods, such as fibrous meats, vegetables, and poultry. It is also preferred that the supplement include between about 100 mg and 200 mg per day of betaine, with about 150 mg per day being most preferred.

Herbs beneficial against the causes of symptoms of diabetes, including high blood pressure, may also be incorporated into the supplement without departing from the scope of the  
20 invention. These herbs include, but are not limited to, ginseng, huckleberry, evening primrose oil, garlic, gotu kola, juniper berries, and suma. The preferred herbs are ginseng and huckleberry. The preferred dose of ginseng is between about 150 mg and about 350 mg per day, with about 275 mg per day being most preferred. The preferred dose of huckleberry is between about 100 mg and about 200 mg per day, with about 150 mg per day being most preferred.

25 Guar gum is a water soluble fiber that has been shown to lower blood glucose levels, aid in lowering cholesterol levels, and curb the appetite. Guar gum must be taken as a liquid as

should not be included in any pills, capsules, tablets or other solid forms. However, the nutritional supplement of the present invention could be prepared in the form of a liquid drink which could include guar gum.

The supplements of the present invention can be manufactured in accordance with procedures known in the art. While the supplement may be formed into a pill with starch or a liquid drink, it is generally preferred that the supplement be prepared in a capsule.

#### EXAMPLE 1

Two nutritional supplements, Phase I and Phase II, were prepared according to the following daily dosages:

##### Phase I

	Vanadyl Sulfate	60 mg
	Chromium (Picolinate)	300 mcg
	Chromium (Glucose Tolerance Factor)	100 mcg
15	L-Carnitine	150 mg
	L-Glutamine	300 mg
	Magnesium	380 mg
	Potassium	114 mg
	Calcium	760 mg
20	Copper	2 mg
	Vitamin A	15,000 IU
	Niacinamide (B-3)	50 mg
	Vitamin (B-1)	6 mg
	Vitamin (B-6)	6 mg
25	Selenium	80 mcg
	Zinc	50 mg

	Pancreatin	100 mg
	Papain	75 mg
	Amylase	100 mg
	Betaine (HCL)	75 mg
5	Lipase	150 mg
	Huckleberry	150 mg
	Ginseng	275 mg

**Phase II**

10	Gymnema sylvestre (extract)	750 mg
	Lipoic acid	100 mg
	Cat's claw	500 mg
	Pullunan	350 mg
15	L-Methionine	200 mg
	Pancreatin	100 mg
	Lipase	100 mg
	Amylase	100 mg
	Dandelion root	300 mg
20	Folic Acid	400 mcg
	Copper (chelated)	2 mg
	Iodine	150 mcg
	Manganese (chelated)	7.5 mg
	Selenium (chelated)	80 mcg
25	Zinc (chelated)	50 mg
	Vitamin B-1	6 mg



Niacinamide (B-3)	50 mg
Vitamin B-6	6 mg

The daily dosages of both the Phase I supplement or the Phase II supplement were divided into six (6) capsules. Two of these six capsules were taken along with breakfast, lunch and dinner each day. Phase I was administered for a period of three months, then Phase II was administered for a period of three months. This pattern of alternating between Phases I and II was repeated to provide continuous benefits to diabetics while avoiding desensitization that can occur from continuous use of a single supplement formulation.

While much of the foregoing disclosure has focused on insulin lack, diabetes can also occur in individuals whose pancreas is producing plenty of insulin but the cells of the body are insulin resistant. Vanadyl sulfate can increase the response that a cell has to insulin by activating the insulin receptor sites on the surface of the cell and within the cell. Furthermore, this activation of insulin receptors may be permanent.

The nutritional supplement of the present invention may be used not only as a treatment for poor glucose metabolism or diabetes, but also for prevention of diabetes by giving the metabolism a boost before full-blown diabetes develops. In fact, because the activation of insulin receptors and other effects may be permanent, the supplement could be considered to be a cure for diabetes in some individuals and circumstances.

It will be understood that certain combinations and sub combinations of the invention are of utility and may be employed without reference to other features in sub combinations. This is contemplated by and is within the scope of the present invention. As many possible embodiments may be made of this invention without departing from the spirit and scope thereof, it is to be understood that all matters hereinabove set forth or shown in the accompanying drawing are to be interpreted as illustrative and not in a limiting sense.

While the foregoing is directed to the preferred embodiment, the scope thereof is determined by the claims which follow:

**WHAT IS CLAIMED IS:**

- 1 1. A nutritional system, comprising:
  - 2 (a) a first supplement comprising a source of vanadate and a source of chromium; and
  - 3 (b) a second supplement comprising *Gymnema sylvestre* and lipoic acid.
- 1 2. The nutritional system of claim 1, wherein the source of vanadate is vanadyl sulfate.
- 1 3. The nutritional system of claim 1, wherein the source of chromium is selected from the  
2 group consisting of chromium picolinate, chromium glucose tolerance factor, and mixtures  
3 thereof.
- 1 4. The nutritional system of claim 4, wherein the source of chromium comprises chromium  
2 picolinate and chromium glucose tolerance factor.
- 1 5. The nutritional system of claim 1, wherein the first supplement further comprises L-  
2 carnitine.
- 1 6. The nutritional system of claim 1, wherein the *Gymnema sylvestre* is provided as an  
2 extract from *Gymnema sylvestre* leaves.
- 1 7. A method of administering nutritional supplements, comprised of:
  - 2 (a) administering a first supplement comprising a source of vanadate; and then
  - 3 (b) administering a second supplement comprising an ingredient selected from *gymnema*  
4 *sylvestre*, lipoic acid or combinations.

1 8. The method of claim 7, further comprising:

2 (c) repeating steps (a) and (b).

1 9. The method of claim 7, wherein the first nutritional supplement further comprises a  
2 source of chromium.

1 10. The method of claim 7, wherein the first nutritional supplement comprises *Gymnema*  
2 *sylvestre* and lipoic acid, and the second nutritional supplement comprises a source of vanadate  
3 and a source of chromium.

1 11. The method of claim 7, wherein the first supplement is administered over a first time  
2 period, the second supplement is administered over a second time period, and the first and  
3 second time periods are different.

1 12. The method of claim 15, wherein the first and second time periods are between about 2  
2 and about 6 months.

1 13. The method of claim 15, wherein the first supplement is administered over a first time  
2 period, the second supplement is administered over a second time period, and the first and  
3 second time periods are about 3 months.

1 14. The method of claim 17, wherein the first and second time periods are 90 days.

1 15. A nutritional system for improving glucose metabolism, comprising:

2 (a) a first supplement comprising a source of vanadate; and

3 (b) a second supplement comprising an ingredient selected from the group consisting

4 of gymnema sylvestre, lipoic acid and combinations thereof.

1 16. The nutritional system of claim 19, wherein the first supplement further comprises a  
2 source of chromium.

1 17. A method of administering nutritional supplements, comprised of:

2 (a) administering a first supplement comprising an ingredient selected from gymnema  
3 sylvestre, lipoic acid and combinations thereof; and then

4 (b) administering a second supplement comprising a source of vanadate.

1 18. The method of claim 21, further comprising:

2 (c) repeating steps (a) and (b).

1 19. The method of claim 21, wherein the second nutritional supplement further comprises a  
2 source of chromium.

1 20. The method of claim 21, wherein the first supplement is administered over a first time  
2 period, the second supplement is administered over a second time period that is different than the  
3 first time period.

1 21. The method of claim 21, wherein the first supplement is administered over a first time  
2 period, the second supplement is administered over a second time period, and the first and  
3 second time periods are between about 2 and about 6 months.

1 22. The method of claim 21, wherein the first supplement is administered over a first time  
2 period, the second supplement is administered over a second time period, and the first and

3 second time periods are about 3 months.

1 23. The method of claim 26, wherein the first and second time periods are 90 days.

1 24. A nutritional supplement for improving glucose metabolism, comprising:

2 (a) a source of vanadate; and

3 (b) a source of chromium.

1 25. The nutritional supplement of claim 28, wherein the source of vanadate is vanadyl  
2 sulfate.

1 26. The nutritional supplement of claim 28, wherein the source of chromium is chromium  
2 picolinate.

1 27. The nutritional supplement of claim 28, further comprising L-carnitine.

1 28. The nutritional supplement of claim 28, further comprising vitamin B1, vitamin B6 and  
2 iron.

1 29. A nutrition supplement for improving glucose metabolism, comprising:

2 (a) gymnema sylvestre; and

3 (b) lipoic acid.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A23L 1/304, 1/30, A61K 33/24</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 98/42211</b> <b>(43) International Publication Date:</b> 1 October 1998 (01.10.98)
<b>(21) International Application Number:</b> PCT/US98/05646 <b>(22) International Filing Date:</b> 24 March 1998 (24.03.98) <b>(30) Priority Data:</b> 08/822,483 24 March 1997 (24.03.97) US <b>(71) Applicant:</b> PROGRESSIVE RESEARCH LABS [US/US]; Suite 162, 9219 Katy Freeway, Houston, TX 77024 (US). <b>(72) Inventor:</b> WOMACK, Rick, W.; Suite 50, 9707 Richmond Avenue, Houston, TX 77042 (US). <b>(74) Agent:</b> STREETS, Jeffrey, L.; Patterson & Streets, L.L.P., Suite 1500, 3040 Post Oak Boulevard, Houston, TX 77056 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 30 December 1998 (30.12.98)
<b>(54) Title:</b> NUTRITIONAL SUPPLEMENTS FOR IMPROVING GLUCOSE METABOLISM  <b>(57) Abstract</b>  The invention provides nutritional supplements and methods for administering nutritional supplements that improve glucose metabolism, particularly for persons with diabetes. A first nutritional supplement, or "Phase I" supplement, comprises a source of vanadate and a source of chromium. A second nutritional supplement, or "Phase II" supplement, comprises <i>Gymnema sylvestre</i> and lipoic acid. The nutritional supplements are alternated to prevent accumulation of the nutrients in the body and also to overcome desensitization that can occur over long periods of continuous use. While the nutritional supplements may be alternated at almost any frequency and taken over almost any duration, it is preferred that each Phase be taken for between about 2 and about 6 months, most preferably about 3 months or about 90 days, before alternating back to the other Phase.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



# INTERNATIONAL SEARCH REPORT

National Application No

PCT/US 98/05646

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A23L1/30 A23L1/30 A61K33/24

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A23L A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 474 358 A (DAI-NIPPON SUGAR) 11 March 1992 see page 2, line 17-21 see page 2, line 35-36 see page 5, line 10-11 see claims; example 1 ---	1-6, 15, 29
A	US 5 116 820 A (Y.HIJI) 26 May 1992  see column 2, line 3-12 see column 2-35-40 see column 2, line 48-53 see claims --- -/--	1-6, 15, 29



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

22 October 1998

Date of mailing of the international search report

29/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van Moer, A

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/05646

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A X	US 5 164 384 A (S.M.PAUL) 17 November 1992 see column 1, line 50 - column 2, line 3 see column 2, line 63 - column 3, line 10 see column 4, line 39-63 see claims ---	1-6,15 24-28
A	WO 94 02036 A (METAGENICS) 3 February 1994  see claims 1,4-7 ---	1-6,15, 24-29
P,A X	US 5 614 224 A (R.W.WOMACK) 25 March 1997 see column 4, line 60-65 see column 5, line 10-25 see column 5, line 34-44 see column 6, line 15-20 see claims -----	1-6,15 24-28

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/05646

## B x I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 7-14, 17-23  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Rule 39.1(iv) PCT- Method for treatment of the human or animal body by therapy
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## B x II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

1st Application No

PCT/US 98/05646

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 474358	A	11-03-1992	JP 7233051 A CA 2048043 A US 5137921 A	05-09-1995 04-03-1992 11-08-1992
US 5116820	A	26-05-1992	JP 1642389 C JP 3008188 B JP 62236469 A	28-02-1992 05-02-1991 16-10-1987
US 5164384	A	17-11-1992	NONE	
WO 9402036	A	03-02-1994	US 5292538 A AU 669003 B AU 4992893 A EP 0651617 A	08-03-1994 23-05-1996 14-02-1994 10-05-1995
US 5614224	A	25-03-1997	US 5730988 A	24-03-1998